

Meteorology 3510
Exercise 11: Due April 22, 2008

This exercise is about microburst downdrafts.

1. Use the same procedure as we used in class to determine the downdraft speed at the surface (where $p = 840$ mb) for a parcel that descends from the environmental LCL (at $p = 590$ mb, $T = 1^\circ\text{C}$) and remains saturated due to rain evaporation until either 1, 2, or 3 g kg^{-1} of rain have evaporated into it, then descends dry adiabatically to the surface. For each case:
 - (a) What is the SEL (sinking evaporation level)?
 - (b) What are the parcel's mixing ratio, T , T_d , and RH at the surface?
 - (c) What is the environment's mixing ratio, T , T_d , and RH at the surface?
 - (d) What is the downdraft CAPE for the parcel?
 - (e) What is the downdraft speed at the surface?

Answers for evaporation of 1 g kg^{-1} of rain: (a) 630 mb, (b) $T = 27^\circ\text{C}$, RH = 29%, (c) $T = 30^\circ\text{C}$, RH = 22%, (d) 230 J kg^{-1} (e) 21 m s^{-1} .
2. For the same environment as Problem 1, the parcel properties at the surface are $T = 24^\circ\text{C}$ and $T_d = 10.5^\circ\text{C}$? For this case,
 - (a) What are the parcel's mixing ratio and RH at the surface?
 - (b) What is the SEL (sinking evaporation level)?
 - (c) How much rain was evaporated into the parcel? (d) What is the downdraft CAPE for the parcel?
 - (e) What is the downdraft speed at the surface?
3. This is like Problem 2, but the environment properties at the surface are $p = 800$ mb, $T = 30^\circ\text{C}$, and $T_d = 3^\circ\text{C}$, and the parcel properties at the surface are $T = 25^\circ\text{C}$ and $T_d = 7^\circ\text{C}$? For this case,
 - (a) What are the parcel's mixing ratio and RH at the surface?
 - (b) What is the SEL (sinking evaporation level)?
 - (c) How much rain was evaporated into the parcel? (d) What is the downdraft CAPE for the parcel?
 - (e) What is the downdraft speed at the surface?