

# Enhanced Mid-Latitude Tropospheric Warming in Satellite Measurements

Qiang Fu,<sup>1,2\*</sup> Celeste M. Johanson,<sup>1</sup> John M. Wallace,<sup>1</sup> Thomas Reichler<sup>3</sup>

**B**oreal summers that follow strong El Niño events, like the one that occurred in 1997, are often characterized by anomalous tropospheric warmth in mid-latitudes of both the Northern and Southern Hemisphere. This warmth occurs in response to El Niño/Southern Oscillation (ENSO)-induced, positive-sea-surface temperature (SST) anomalies in the Indian and western Pacific Ocean “warm pool” regions (1). The anomalous mid-latitude tropospheric warmth is accompanied by an elevation of the pressure surfaces in the upper troposphere and an increased frequency of droughts, including in parts of the United States (1, 2).

From 1979 onward, the most pronounced SST warming has occurred within this same “warm pool” region (3). If the planetary-scale atmospheric circulation responds to the multidecadal SST trends in the same way as it responds to ENSO-induced SST variability, one might expect to observe a multidecadal trend toward a warmer mid-latitude troposphere.

We examined atmospheric temperature trends for 1979 to 2005 based on satellite-borne microwave sounding unit (MSU) data (4–6). Figure 1 shows the global spatial patterns of strato-

spheric and tropospheric temperature trends for 1979 to 2005 and the corresponding zonal mean latitudinal profiles. To emphasize the spatial gradients in the trends, the reference (white) values in the color bars in (Fig. 1, A and B) are set equal to the respective global mean trends. The most pronounced feature is the enhanced stratospheric cooling and tropospheric warming in the 15 to 45° latitude belts in both hemispheres, relative to other latitudes. The atmospheric trends for 1979 to 1997 (fig. S1) exhibit features similar to those in Fig. 1, which indicates that existence of the trend is not contingent on the episode of record-high mid-latitude temperatures that occurred in summer 1998 in response to the 1997 El Niño (1) but is a robust feature of the period of record from 1979 onward (fig. S2). The same pattern is evident in the trend in radiances from MSU channel 2 (fig. S3), which is a further proof of the enhanced mid-latitude tropospheric warming corresponding to the enhanced stratospheric cooling.

Unless it is compensated by a drop in sea-level pressure in the same latitude belt, the observed pattern of temperature changes in Fig. 1 is indicative of a tendency toward an upward bulging of the upper tropospheric pressure surfaces centered around 30° latitude in both hemispheres. Because 30° latitude corresponds to the latitude of the tropospheric jet streams, it can be inferred that the downward slope of the pressure surfaces toward the poles has been increasing on the poleward flanks of the jet streams and decreasing on the equatorward flanks. Such a reshaping of the pressure surfaces would have the effect of shifting the jet streams poleward. Based on the observed temperature changes alone, we estimate that the jet streams in both hemispheres have shifted poleward by  $\sim 1^\circ$  latitude in both summer and winter seasons (6). Because the jet streams mark the poleward limit of the tropical Hadley circulation,

a systematic poleward shift of the jet streams implies that the tropical circulation has widened by  $\sim 2^\circ$  latitude during this 27-year period (7). (Our analysis of the National Centers for Environmental Prediction/National Center for Atmospheric Research reanalyses suggests that sea-level pressures near 30°N and 30°S have risen relative to surrounding latitudes. Such pressure rises, if real, would cause an additional poleward shift in the jet streams.)

In contrast to the seasonally dependent circulation changes reported in association with the Northern and Southern Hemisphere annular modes, the changes reported here are occurring at somewhat lower latitudes, and the Northern Hemisphere trends are no less pronounced during the warm season (fig. S4) than during the cold season.

Whether the observed trends in Fig. 1 are an integral part of the response to greenhouse warming remains to be seen (8–10). Regardless of the cause, the poleward shift of the jet streams and the associated subtropical dry zone (11), if it continues, could have important societal implications.

## References and Notes

- N. C. Lau, A. Leetmaa, M. J. Nath, H. L. Wang, *J. Clim.* **18**, 2922 (2005).
- M. P. Hoerling, A. Kumar, *Science* **299**, 691 (2003).
- T. M. Smith, R. W. Reynolds, *J. Clim.* **18**, 2021 (2005).
- C. A. Mears, M. C. Schabel, F. J. Wentz, *J. Clim.* **16**, 3650 (2003).
- Q. Fu, C. M. Johanson, S. G. Warren, D. J. Seidel, *Nature* **429**, 55 (2004).
- Materials and methods are available as supporting material on Science Online.
- T. Reichler, I. Held, paper presented at the 17th Conference on Climate Variability and Change, Cambridge, MA, 13 to 17 June 2005.
- State-of-the-art coupled ocean-atmospheric models predict stronger warming in the tropical troposphere than in mid-latitudes (9) and a poleward shift of the eddy-driven jets centered  $\sim 45^\circ$  latitude (10). A pattern like the recent trends, with strongest tropospheric warming centered  $\sim 30^\circ$  latitude, is not recovered in the simulations.
- J. T. Houghton, *Climate Change 2001: The Scientific Basis* (Cambridge Univ. Press, Cambridge, 2001).
- S. S. Leroy, J. G. Anderson, J. A. Dykema, paper presented at the 18th Conference on Climate Variability and Change, Atlanta, GA, 28 January to 3 February 2006.
- S. D. Schubert *et al.*, *J. Clim.* **17**, 485 (2004).
- This study is supported by the National Oceanic and Atmospheric Administration grant NA17RJ1232 and NASA grant NNG04GM23G. We thank D. L. Hartmann and J. Paele for useful discussions.


## Supporting Online Material

www.sciencemag.org/cgi/content/full/312/5777/1179/DC1  
Materials and Methods  
Figs. S1 to S4  
References

30 January 2006; accepted 30 March 2006  
10.1126/science.1125566

<sup>1</sup>Department of Atmospheric Sciences, University of Washington, Seattle, WA 98195, USA. <sup>2</sup>College of Atmospheric Sciences, Lanzhou University, Lanzhou, Gansu, 730000, China. <sup>3</sup>Department of Meteorology, University of Utah, 135 S 1460 E, Room 819 (WBB), Salt Lake City, UT 84112-0110, USA.

\*To whom correspondence should be addressed. E-mail: qfu@atmos.washington.edu



**Fig. 1.** Global and latitudinal distributions of atmospheric temperature trends for 1979 to 2005 based on satellite-borne MSU observations. (A and C) Stratospheric trends with a global mean of  $-0.33$  K per decade. (B and D) Tropospheric trends with a global mean of  $+0.20$  K per decade. Orange shading represents enhanced rates of stratospheric cooling and tropospheric warming relative to their respective global means, and blue shading represents suppressed rates. The polarity is reversed to facilitate comparison between (A) and (B).