Lake- and Sea-Effect Precipitation

VU2: Course Number 707813

Jim Steenburgh
Fulbright Visiting Professor of Natural Sciences
University of Innsbruck
Department of Atmospheric Sciences
University of Utah
jim.steenburgh@utah.edu

Learning Objectives

- After this class you should be able to
  - Recognize several ways that lakes and complex terrain affect the morphology and intensity of lake-effect storms
  - Use this recognition to better analyze and predict lake-effect storms

Discussion

What is lake- and sea-effect precipitation/snow?

What Is Lake- and Sea-Effect Precip?

- "Lake effect snow i[...] produced w[...]en cold winds move across long [...]xpa[...]es [...] wa[...]mer lake water, *picking up water* [...]*apor,* w[...]ch fre[...]zes and is deposited on the [...]e sh[...]e."
  – W[...]kipedia.c[...]n (2006)
- "Precipitation occurring near or downwind from the shore of a lake resulting from the *warming (destabilization) and moistening* of relatively cold air during passage over a warm body of water
  – Glossary of Meteorology (2000)

Lake- and Sea-Effect Precipitation



Courtesy Peter Veals

Precipitation produced primarily by boundary layer convection that is generated, enhanced, and organized by sensible and latent heat fluxes and associated boundary layer and mesoscale circulations as cold air moves over relatively warm water

Key Ingredients

- Instability
  - Lake–850-hPa or Lake–700-hPa ΔT exceeding dry adiabatic
    - There are some exceptions
- Wind Direction/Fetch
- Moisture if fetch is small can be important
  - Less important for large bodies of water
- Boundary layer or mesoscale circulations
  - Former include land breezes and terrain-forced flows

Steenburgh et al. (2000), Alcott et al. (2012)

Additional Factors

- Shoreline geometry
- Land and lake breezes
- Upstream and downstream topography
- Multi-lake/sea effects
- Ice cover (where and when it happens)

Example



Lake- and Sea-Effect Regions



Sea of Japan, Yellow Sea, Sea of Okhotsk

Lake- and Sea-Effect Regions



Baltic Sea, North Sea, English Channel, Norwegian Sea

Lake- and Sea-Effect Regions



Eastern Mediterranean, Black Sea

Lake- and Sea-Effect Regions



Central Mediterranean, Adriatic Sea

Lake- and Sea-Effect Regions



North American
Great Lakes

"Small" Lakes



Lake Constance, CH/DE/AT



Great Salt Lake, USA



Finger Lakes, USA

Umek and Gohm 2016; Alcott et al. 2012; Bergmaier and Geerts 2016

Extremes



Lake Ontario & Tug Hill Plateau
Elongated body of water 300 km long
Tug Hill Plateau rises 500 m above lake level

Sea of Japan and Japanese Mountains
Large body of water with NW fetch ~850 km
Downstream terrain rises ~1000–2800 m

Tug Hill: Storms of Great Intensity

"Snow rates during some events are the greatest ever measured on record from anywhere in the world"
– Burt (2007)

35 cm in 1 hour
Copenhagen, NY
2 Dec 1966

130 cm in 16 hours
Bennetts Bridge, NY
17-18 Jan 1958

196 cm in 24 hours*
Montague, NY
11-12 Jan 1997

*Based on 6 measurements

Overnight Snowfall, Adams, New York, USA
OWLeS IOP7

Burt (2007)

Tug Hill Maximum



Leffler et al. (1997)

Japan's Gosetsu Chitai (Heavy Snow Region)



Sapporo
Pop: 1.95 million
Mean Annual Snowfall: 630 cm

Niseko United Resorts
Mean Annual Snowfall: 1400 cm

Sukayu Onsen
Mean Annual Snowfall: 1764 cm

Tsunan
Mean Annual Snowfall: 1349 cm

Japan's Gosetsu Chitai (Heavy Snow Region)

Photo: J. Steenburgh

"The surest climatological bet for deep powder skiing anywhere in the world"

Photo: Tateyama-Kurobe Snow Corridor



"Some of the world's deepest seasonal snowpacks"- Yamaguchi et al. (2011)

Morphology

Lake- and sea-effect modes

- "Broad coverage"
  - Open cells/non-banded
  - Longitudinal-mode bands
  - Transverse-mode bands
- Mesoscale bands
  - Long-lake-axis parallel (LLAP)
  - Japan Sea Polar Airmass Convergence Zone
  - Other terrain/coastally forced
- Mesovorticies

Open Cells/Non-Banded





Alcott et al. 2012; Campbell et al. 2016

Longitudinal Mode (Cloud Streets)



http://www.eumetrain.org/satmanu/CMs/ClStr/print.htm

Horizontal Roll Convection

Formed by buoyancy and shear

Nearly parallel to ambient flow

Transverse Mode





Horizontal Roll Convection

High directional shear environments

Parallel to shear vector

Nearly transverse to mean flow

Asai 1972; Campbell et al. 2018

LLAP Bands (Type I)







Flow along major lake axis

Large lake-land ΔT

Mesoscale forcing from land breeze or breezes

Paul Markowski; Alcott et al. 2012; Campbell et al. 2016

LLAP Band (Type I) Event



LLAP Bands (Type II)





Form near lee shore

Land breeze opposes large-scale flow

Veals and Steenburgh 2015; Hjelmfelt 1990

JPCZ



Japan Sea Polar Airmass Convergence Zone

Drivers include convergence in lee of Korean Highlands, coastal geometry, Sea of Japan SST distribution

Courtesy Tyler West

Other Terrain/Coastal Effects



Mesovortex



https://epod.usra.edu/blog/2008/06/lake-michigan-mesovortex-radar.html

Discussion

What controls the mode of lake- and sea-effect precipitation?

Ontario Winter Lake-effect Systems (OWLeS)Field Campaign

OWLeS Orographic Transect



Prior Conceptual Model



Lackmann (2011), Minder et al. (2015)

Tug Hill: Inland/Orographic Transition





Instead: Convective to stratiform transition with snowfall
Becoming more continuous and persistent

Minder et al. (2015)

Mesoscale & Orographic Forcing OWLeS IOP2b

0000/11–0000/12 UTC Dec





Long-Lake Axis Parallel (LLAP) System

SC
Snow: 48 cm
LPE: 33.5 mm

NR
Snow: 102 cm
LPE: 62.5 mm

OR=NR/SC=1.9

Campbell et al. (2016)

Forcing Conceptual Models



Thermally Forced Land Breeze Convergence
(Symmetric, Mid-Lake Axis)

UCAR/Comet

Forcing Conceptual Models



Frictionally Forced Land Breeze Convergence
(Streamwise-Right Shore)

Markowski and Richardson (2010)

Lake Ontario Not Symmetrical Or Oval!



Impact of Shoreline Geometry

1200 UTC 11 Jan: WRF near-surface wind and divergence ($x10^{-4}$ s^{-1})



Steenburgh and Campbell (2017)

Impact of Shoreline Geometry

1200 UTC 11 Jan: WRF near-surface pot. temp. and pot. temp. gradient (K/10 km)



Steenburgh and Campbell (2017)

Impact of Shoreline Geometry

1200 UTC 11 Jan: WRF near-surface wind, IR, reflectivity, potential temperature



Steenburgh and Campbell (2017)

Impact of Shoreline Geometry



- Cross west shore, long over-lake residence time
- Cross north or south shore, short over-lake residence time
- Confined to land, little or no lake modification

Steenburgh and Campbell (2017)

Getting Back to Tug Hill

1700 UTC 11 Jan: WRF theta-e (black contours), ascent (red contours), section-parallel circulation vectors, and hydrometeor mixing ratio (shaded)





Zonal cross section across Tug Hill Plateau

Campbell and Steenburgh (2017)

Getting Back to Tug Hill

1930 UTC 11 December





Bergmaier et al. (2017)

UWKA Cloud Radar

1928–1935 UTC 11 December





Bergmaier et al. (2017)

Model Terrain Sensitivity



Campbell and Steenburgh (2017)

Tug Hill Influences



0300-2200 UTC
Vertical Velocity

0300-2200 UTC
Precipitation

Campbell and Steenburgh (2017)

Tug Hill Influences



Watson and Lane (2012)

Japan’s Gosetsu Chitai



Yamayuki or Satoyuki?



Yamayuki
Mountain Snowfall

Satoyuki
Lowland Snowfall



Conceptual Model of Satoyuki Snowfall
Magono et al. (1966)

Discussion

What controls whether a storm is Yamayuki or Satoyuki?

Hokuriku Region



Wind direction (Moderate *U*)



Veals et al. (2019, submitted)

Wind Speed (290˚)



Veals et al. (2019, submitted)

Tug Hill Comparison



Veals et al. (2018)

Wind Speed (320˚)



Veals et al. (2019, submitted)

Non-Dimensional Mountain Height



Veals et al. (2019, submitted)

Lake Constance



Observed Event



0800 UTC 8 Feb 2013

Precip
Radar: 0640–1200 UTC 8 Feb
Gauge: 0700–1200 UTC 8 Feb

Umek and Gohm 2016

Observed Event



Importance of blocking by topography south of Lake Constance

Umek and Gohm 2016

Simulation



Importance of blocking by topography south of Lake Constance

Umek and Gohm 2016

Sensitivity Studies



Control

No terrain

No terrain to SE

Conclusion: Terrain blocking critical

Umek and Gohm 2016

Cloud Modeling



Oval Lake LLAP Band



Courtesy Tom Gowan

Broad Lake Open Cellular



Courtesy Tom Gowan

Oval vs. Wide/LLAP vs. Open Cell



Discussion: With same upstream sounding and terrain, why is the orographic ratio different in these two cases?

Courtesy Tom Gowan

Summary

- Many processes influence lake-effect systems
  - Upstream instability and moisture
  - Lake conditions (surface temperature, sub-surface temperature, salinity, ice cover)
  - Land breezes and PBL circulations
  - Orography
- Orographic influences are multifaceted and affect the initiation, intensity, and morphology of lake-effect systems

References

Alcott, T. I., W. J. Steenburgh, and N. F. Laird, 2012: Great Salt Lake-effect precipitation: Observed frequency, characteristics, and environmental factors. *Wea. Forecasting*, **27**, 954-971.

Asai, T., 1972: Thermal instability of a shear flow turning the direction with height. *J. Meteor. Soc. Japan*, **50**, 525–532

Bergmaier, P.T. and B. Geerts, 2016: Airborne radar observations of lake-effect snowbands over the New York Finger Lakes. *Mon. Wea. Rev*., **144**, 3895–3914.

Bergmaier, P. T., B. Geerts, L. S. Campbell, and W. J. Steenburgh, 2017: The OWLeS IOP2b lake-effect snowstorm: Dynamics of the secondary circulation. *Mon. Wea. Rev*., **145**, 2437-2459.

Burt, C. C., 2007: *Extreme Weather: A Guide and Record Book*. W. W. Norton Company, 320 pp.

Campbell, L. S., and W. J. Steenburgh, 2017: The OWLeS IOP2b lake-effect snowstorm: Mechanisms contributing to the Tug Hill Precipitation Maximum. *Mon. Wea. Rev*., **145**, 2461-2478.

Campbell, L. S., W. J. Steenburgh, P. G. Veals, T. W. Letcher, and J. R. Minder, 2016: Lake-effect mode and precipitation enhancement over the Tug Hill Plateau during OWLeS IOP2b. *Mon. Wea. Rev*., 144, 1729–1748.

Campbell, L. S., W. J. Steenburgh, Y. Yamada, M. Kawashima, and Y. Fujiyoshi, 2018: Influences of orography and coastal geometry on a transverse-mode sea-effect snowstorm over Hokkaido Island, Japan. *Mon. Wea. Rev*., **146**, 2201–2220.

Hjelmfelt, M. R., 1990: Numerical study of the influence of environmental conditions on lake-effect snowstorms over Lake Michigan. *Mon. Wea. Rev*., **118,** 138-150.

Lackmann, G., 2011: *Midlatitude Synoptic Meteorology*. Amer. Meteor. Soc., 360 pp.

Leffler, R. J., R. M. Downs, G. W. Goodge, N. J. Doesken, K. L. Eggleston, and D. Robinson, 1997: Evaluation of the reported January 11–12, 1997, Montague, New York, 77-inch, 24-hour lake-effect snowfall. National Weather Service Special Rep., 60 pp. [Available online at http://www1.ncdc.noaa.gov/pub/data/cmb/extremes/ncec/mantague-ny-snowfall-24hour.pdf.]

References

Magono, C., K. K. Kikuchi, T. Kimura, S. Tazawa, and T. Kasai, 1966: A study of the snowfall in the winter monsoon season in Hokkaido with special reference to low land snowfall. *J. Fac. Sci., Hokkaido Univ.*, **11**, 287-308.

Markowski, P., and Y. Richardson, 2010: *Mesoscale Meteorology in Midlatitudes*. Wiley-Blackwell, 407 pp.

Minder, J. R., T. Letcher, L. S. Campbell, P. G. Veals, and W. J. Steenburgh, 2015: The evolution of lake-effect convection during landfall and orogrpahic uplift as observed by profiling radars. *Mon. Wea. Rev*., **143**, 4422–4442.

Umek, L. and A. Gohm, 2016: Lake and orographic effects on a snowstorm at Lake Constance. *Mon. Wea. Rev*., **144**, 4687–4707

Steenburgh, W. J., and L. S. Campbell, 2017: The OWLeS IOP2b lake-effect snowstorm: Shoreline geometry, airmass boundaries, and the mesoscale forcing of precipitation. *Mon. Wea. Rev*., **145**, 2421-2436.

Steenburgh, W. J., S. F. Halvorson, and D. J. Onton, 2000: Climatology of lake-effect snowstorms of the Great Salt Lake. *Mon. Wea. Rev*., **128,** 709-727.

Veals, P. G., and W. J. Steenburgh, 2015: Climatological characteristics and orographic enhancement of lake-effect precipitation east of Lake Ontario and over the Tug Hill Plateau. *Mon. Wea. Rev*., **143**, 3591-3609.

Veals, P. G., W. J. Steenburgh, and L. S. Campbell, 2018: Factors affecting the inland and orographic enhancement of lake-effect precipitation over the Tug Hill Plateau. *Mon. Wea. Rev*., **146**, 1745–1762.

Veals, P. G., W. J. Steenburgh, S. Nakai, and S. Yamaguchi, 2019: Factors affecting the inland and orographic enhancement of sea-effect snowfall in the Hokuriku Region of Japan. Submitted to *Mon. Wea. Rev*.

Watson, C. D., and T. P. Lane, 2012: Sensitivities of orographic precipitation to terrain geometry and upstream conditions in idealized simulations. *J. Atmos. Sci*., **69**, 1208–1231.